## Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2016 (Page 1)

Vaccine	Child Care Facility,	Kindergarten through 12 <sup>th</sup> Grade		Minimum Intervals
Requirement	Preschool, Early Childhood Pre-Kindergarten Programs	First Entry into School (Kindergarten or First Grade)	Other Grades	Allowed Between Doses and Other Options for Proof of Immunity
DTP/DTaP/ or Tdap, Td (Diphtheria, Tetanus, Pertussis)	Three doses by 1 year of age  One additional booster dose by 2 <sup>nd</sup> birthday	Four or more doses of DTP/DTaP with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday	Three or more doses of DTP/DTaP or Td; with the last dose qualifying as a booster if received on or after the 4 <sup>th</sup> birthday  For Students entering 6 <sup>th</sup> thru 12 <sup>th</sup>	Minimum interval between series doses: 4 weeks (28 days) Between series and booster: 6 months
			grades: 1 dose of Tdap	No proof of immunity allowed
Polio	Two doses by 1 year of age.  One additional dose by  2 <sup>nd</sup> birthday	Four or more doses of the same type of Polio vaccine with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday.	Three or more doses of Polio with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday.  *If the series is given in any combination of polio vaccine types, 4 or more doses are required with the last being a	Minimum interval between series doses: 4 weeks (28 days) For Grade K: 6 month interval between 3 dose series and booster; booster must be on or after 4 <sup>th</sup> birthday
Measles	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	(progressive requirement) booster on or after the 4 <sup>th</sup> birthday.  Two doses of Measles Vaccine, the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		No proof of immunity allowed  Laboratory evidence of measles immunity <u>OR</u> Certified physician verification* of measles disease by date of illness *Cases diagnosed after 7/1/2002 must include lab evidence of infection
Rubella	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	<b>Two doses of Rubella Vaccine</b> , the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		Laboratory evidence of rubella immunity History of disease is not acceptable proof of immunity to rubella
Mumps	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	<b>Two doses of Mumps Vaccine</b> , the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		Laboratory evidence of mumps immunity OR Certified physician verification of mumps disease by date of illness
Haemophilus influenzae type b	Refer to Hib ACIP recs for schedule for series Children 24-59 mos. without series must have one dose	Not required after the 5 <sup>th</sup> birthday (60 months of age)		Refer to ACIP Hib vaccination schedule
Invasive Pneumococcal Disease	after 15 mos. of age  Refer to PCV ACIP recs for schedule for series  Children 24-59 mos. without series must have one dose	Not required after the 5 <sup>th</sup> birthday (60 months of age)		No proof of immunity allowed  Refer to ACIP PCV vaccination schedule
	after 24 months of age			No proof of immunity allowed

## Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2016 (Page 2)

Vaccine	Child Care Facility, Preschool, Early Childhood Pre-Kindergarten Programs	Kindergarten through 12 <sup>th</sup> Grade		Minimum Intervals
Requirement		First Entry into School (Kindergarten or First Grade)	Other Grades	Allowed Between Doses and Other Options for Proof of Immunity
Hepatitis B	Three doses for all children  Third dose must have been administered on or after 6 months of age (168 days)	No Requirements	For Students entering grades 6 thru 12:  Three doses hepatitis B vaccine administered at recommended intervals	Minimum intervals between doses:  1 & 2- at least 4 weeks (28 days)  2 & 3 - at least 2 months (56 days)  1 & 3 - at least 4 months (112 days)  Laboratory evidence of prior or current infection
Varicella (progressive requirement)	One dose on or after 1 <sup>st</sup> birthday	Two doses of Varicella Vaccine, for Students entering Kindergarten and grades and 1  The 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.	One dose of Varicella on or after the 1 <sup>st</sup> birthday for Students entering grades 1 thru 12	Minimum intervals for administration: The 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.  Statement from physician or health care provider verifying disease history OR  Laboratory evidence of varicella immunity
			Two doses of Varicella Vaccine for Students entering grades 2, 6, 7, 8, 9, 10, & 11	
Meningococcal Conjugate Vaccine (progressive requirement)	No Requirements	No Requirements	Applies to Students entering grades 6, 7, or 12 beginning 2016-2017 school year  1 dose of Meningococcal vaccine for entry to grade 6 or 7  2 doses of Meningococcal vaccine at	Minimum intervals for administration: The first dose received on or after the 10 <sup>th</sup> birthday; second dose at least eight weeks after the 1 <sup>st</sup> dose. Only one dose is required if the first dose was received at 16 years
			entry to 12 <sup>TH</sup> grade	of age or older.  No proof of immunity allowed.

Source: Child Health Examination Code/Part 665,
Prepared by: Illinois Department of Public Health Immunization Section, January 2016